Electronic Journal: Southwest Journal of Pure and Applied Mathematics
Internet: http://rattler.cameron.edu/swjpam.html
ISSN 1083-0464
Issue 2, December 2000, pp. 152–159.
Submitted: September 13, 2000. Published: December 15, 2000.

ABOUT THE EXISTENCE OF THE THERMODYNAMIC LIMIT FOR SOME DETERMINISTIC SEQUENCES OF THE UNIT CIRCLE

Stefano Siboni

Abstract. We show that in the set $\Omega = \mathbb{R}_+ \times (1, +\infty) \subset \mathbb{R}^2_+$, endowed with the usual Lebesgue measure, for almost all $(h, \lambda) \in \Omega$ the limit $\lim_{n\to+\infty} \frac{1}{n} \ln \left| h\left(\lambda^n - \lambda^{-n}\right) \mathrm{mod}\, [-1/2, 1/2) \right|$ exists and is equal to zero. The result is related to a characterization of relaxation to equilibrium in mixing automorphisms of the two-torus. It is nothing but a curiosity, but maybe you will find it nice.

1991 A.M.S. Subject Classification Codes:. 28,30,26.

Keywords and Phrases:. Thermodynamic Limit

Dipartimento di Ingegneria dei Materiali Facoltà di Ingegneria, Università di Trento
Trento - Italy
E-mail: Stefano.Siboni@ing.unitn.it
©2000 Cameron University

Typeset by $\mathcal{AMS}$-TEX