

# SLqPCR: Functions for analysis of real-time quantitative PCR data at SIRS-Lab GmbH

Dr. Matthias Kohl  
SIRS-Lab GmbH (Jena, Germany)



May 12, 2008

## Contents

|          |                                                              |          |
|----------|--------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                          | <b>1</b> |
| <b>2</b> | <b>Selection of most stable reference/housekeeping genes</b> | <b>1</b> |
| <b>3</b> | <b>Normalization by geometric averaging</b>                  | <b>7</b> |

## 1 Introduction

The package "SLqPCR" was designed for the analysis of real-time quantitative RT-PCR data. In this short vignette we describe and demonstrate the available functions.

## 2 Selection of most stable reference/housekeeping genes

We describe the selection of the best (most stable) reference/housekeeping genes using method and data set of Vandesompele et al (2002) [1] (in the sequel: Vand02). We load library and data

```
> library(SLqPCR)
> data(vandesompele)
> str(vandesompele)
```

```
'data.frame':      85 obs. of  10 variables:
 $ ACTB   : num  0.0425 0.0192 0.1631 0.5726 0.0370 ...
```

```

$ B2M    : num  0.0576 0.0194 0.2956 1.0000 0.0444 ...
$ GAPD   : num  0.1547 0.0703 0.7733 1.0000 0.1192 ...
$ HMBS   : num  0.110 0.088 0.405 0.797 0.208 ...
$ HPRT1  : num  0.1180 0.0708 0.5575 1.0000 0.1304 ...
$ RPL13A : num  0.0742 0.0441 0.3481 0.5707 0.1078 ...
$ SDHA   : num  0.203 0.140 0.447 0.974 0.214 ...
$ TBP    : num  0.190 0.106 0.469 1.000 0.201 ...
$ UBC    : num  0.0992 0.0368 0.3401 0.5980 0.0759 ...
$ YWHAZ  : num  0.1032 0.0393 0.3588 0.7863 0.1002 ...

```

We start by ranking the selected reference/housekeeping genes. The function `selectHKgenes` proceeds stepwise; confer Section “Materials and methods” in Vand02. That is, the gene stability measure  $M$  of all candidate genes is computed and the gene with the highest  $M$  value is excluded. Then, the gene stability measure  $M$  for the remaining gene is calculated and so on. This procedure is repeated until two respectively `minNrHK` is reached.

```

> tissue <- as.factor(c(rep("BM", 9), rep("POOL", 9), rep("FIB",
+   20), rep("LEU", 13), rep("NB", 34)))
> res.BM <- selectHKgenes(vandesompele[tissue == "BM", ], method = "Vandesompele",
+   geneSymbol = names(vandesompele), minNrHK = 2, trace = TRUE,
+   na.rm = TRUE)

```

```
#####
```

Step 1 :

gene expression stability values  $M$ :

| HPRT1     | YWHAZ     | RPL13A    | UBC       | GAPD      | SDHA      | TBP       | HMBS      |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 0.5160313 | 0.5314564 | 0.5335963 | 0.5700961 | 0.6064919 | 0.6201470 | 0.6397969 | 0.7206013 |
| B2M       | ACTB      |           |           |           |           |           |           |
| 0.7747634 | 0.8498739 |           |           |           |           |           |           |

average expression stability  $M$ : 0.6362855

gene with lowest stability (largest  $M$  value): ACTB

Pairwise variation, ( 9 / 10 ): 0.076469

```
#####
```

Step 2 :

gene expression stability values  $M$ :

| HPRT1     | RPL13A    | YWHAZ     | UBC       | GAPD      | SDHA      | TBP       | HMBS      |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 0.4705664 | 0.5141375 | 0.5271169 | 0.5554718 | 0.5575295 | 0.5738460 | 0.6042110 | 0.6759176 |
| B2M       |           |           |           |           |           |           |           |
| 0.7671985 |           |           |           |           |           |           |           |

average expression stability  $M$ : 0.5828883

gene with lowest stability (largest  $M$  value): B2M

Pairwise variation, ( 8 / 9 ): 0.07765343

```

#####
Step 3 :
gene expression stability values M:
      HPRT1      RPL13A      SDHA      YWHAZ      UBC      GAPD      TBP      HMBS
0.4391222 0.4733732 0.5243665 0.5253471 0.5403137 0.5560120 0.5622094 0.6210820
average expression stability M:          0.5302283
gene with lowest stability (largest M value):          HMBS
Pairwise variation, ( 7 / 8 ):          0.067112
#####

Step 4 :
gene expression stability values M:
      HPRT1      RPL13A      YWHAZ      UBC      SDHA      GAPD      TBP
0.4389069 0.4696398 0.4879728 0.5043292 0.5178634 0.5245346 0.5563591
average expression stability M:          0.4999437
gene with lowest stability (largest M value):          TBP
Pairwise variation, ( 6 / 7 ):          0.06813202
#####

Step 5 :
gene expression stability values M:
      HPRT1      RPL13A      UBC      YWHAZ      GAPD      SDHA
0.4292808 0.4447874 0.4594181 0.4728920 0.5012107 0.5566762
average expression stability M:          0.4773775
gene with lowest stability (largest M value):          SDHA
Pairwise variation, ( 5 / 6 ):          0.08061944
#####

Step 6 :
gene expression stability values M:
      UBC      RPL13A      HPRT1      YWHAZ      GAPD
0.4195958 0.4204997 0.4219179 0.4424631 0.4841646
average expression stability M:          0.4377282
gene with lowest stability (largest M value):          GAPD
Pairwise variation, ( 4 / 5 ):          0.08416531
#####

Step 7 :
gene expression stability values M:
      RPL13A      UBC      YWHAZ      HPRT1
0.3699163 0.3978736 0.4173706 0.4419220
average expression stability M:          0.4067706
gene with lowest stability (largest M value):          HPRT1
Pairwise variation, ( 3 / 4 ):          0.09767827
#####

```

Step 8 :

gene expression stability values M:

UBC RPL13A YWHAZ  
0.3559286 0.3761358 0.3827933

average expression stability M: 0.3716192

gene with lowest stability (largest M value): YWHAZ

Pairwise variation, ( 2 / 3 ): 0.1137450

#####

Step 9 :

gene expression stability values M:

RPL13A UBC  
0.3492712 0.3492712

average expression stability M: 0.3492712

```
> res.POOL <- selectHKgenes(vandesompele[tissue == "POOL", ], method = "Vandesompele",
+   geneSymbol = names(vandesompele), minNrHK = 2, trace = FALSE,
+   na.rm = TRUE)
> res.FIB <- selectHKgenes(vandesompele[tissue == "FIB", ], method = "Vandesompele",
+   geneSymbol = names(vandesompele), minNrHK = 2, trace = FALSE,
+   na.rm = TRUE)
> res.LEU <- selectHKgenes(vandesompele[tissue == "LEU", ], method = "Vandesompele",
+   geneSymbol = names(vandesompele), minNrHK = 2, trace = FALSE,
+   na.rm = TRUE)
> res.NB <- selectHKgenes(vandesompele[tissue == "NB", ], method = "Vandesompele",
+   geneSymbol = names(vandesompele), minNrHK = 2, trace = FALSE,
+   na.rm = TRUE)
```

We obtain the following ranking of genes (cf. Table 3 in Vand02)

```
> ranks <- data.frame(c(1, 1:9), res.BM$ranking, res.POOL$ranking,
+   res.FIB$ranking, res.LEU$ranking, res.NB$ranking)
> names(ranks) <- c("rank", "BM", "POOL", "FIB", "LEU", "NB")
> ranks
```

|   | rank | BM     | POOL   | FIB   | LEU    | NB    |
|---|------|--------|--------|-------|--------|-------|
| 1 | 1    | RPL13A | GAPD   | GAPD  | UBC    | GAPD  |
| 2 | 1    | UBC    | SDHA   | HPRT1 | YWHAZ  | HPRT1 |
| 3 | 2    | YWHAZ  | HMBS   | YWHAZ | B2M    | SDHA  |
| 4 | 3    | HPRT1  | HPRT1  | UBC   | GAPD   | UBC   |
| 5 | 4    | GAPD   | TBP    | ACTB  | RPL13A | HMBS  |
| 6 | 5    | SDHA   | UBC    | TBP   | TBP    | YWHAZ |
| 7 | 6    | TBP    | RPL13A | SDHA  | SDHA   | TBP   |

|    |   |      |       |        |       |        |
|----|---|------|-------|--------|-------|--------|
| 8  | 7 | HMBS | YWHAZ | RPL13A | HPRT1 | ACTB   |
| 9  | 8 | B2M  | ACTB  | B2M    | HMBS  | RPL13A |
| 10 | 9 | ACTB | B2M   | HMBS   | ACTB  | B2M    |


**Remark 1:**

- (a) Since the computation is based on gene ratios, the two most stable control genes in each cell type cannot be ranked.
- (b) In praxis the selection of reference/housekeeping genes may require an additional step which is the computation of relative quantities via `relQuantPCR`; e.g.

```
> exa1 <- apply(vandesompele[tissue == "BM", ], 2, relQuantPCR,
+               E = 2)
```


We plot the average expression stability M for each cell type (cf. Figure 2 in Vand02).

```
> library(RColorBrewer)
> mypalette <- brewer.pal(5, "Set1")
> matplot(cbind(res.BM$meanM, res.POOL$meanM, res.FIB$meanM, res.LEU$meanM,
+               res.NB$meanM), type = "b", ylab = "Average expression stability M",
+               xlab = "Number of remaining control genes", axes = FALSE,
+               pch = 19, col = mypalette, ylim = c(0.2, 1.22), lty = 1,
+               lwd = 2, main = "Gene stability measure")
> axis(1, at = 1:9, labels = as.character(10:2))
> axis(2, at = seq(0.2, 1.2, by = 0.2), labels = as.character(seq(0.2,
+               1.2, by = 0.2)))
> box()
> abline(h = seq(0.2, 1.2, by = 0.2), lty = 2, lwd = 1, col = "grey")
> legend("topright", legend = c("BM", "POOL", "FIB", "LEU", "NB"),
+               fill = mypalette)
```



Second, we plot the pairwise variation for each cell type (cf. Figure 3 (a) in Vand02)

```
> mypalette <- brewer.pal(8, "YlGnBu")
> barplot(cbind(res.BM$variation, res.POOL$variation, res.FIB$variation,
+   res.LEU$variation, res.NB$variation), beside = TRUE, col = mypalette,
+   space = c(0, 2), names.arg = c("BM", "POOL", "FIB", "LEU",
+   "NB"))
> legend("topright", legend = c("V9/10", "V8/9", "V7/8", "V6/7",
+   "V5/6", "V4/5", "V3/4", "V2/3"), fill = mypalette, ncol = 2)
> abline(h = seq(0.05, 0.25, by = 0.05), lty = 2, col = "grey")
> abline(h = 0.15, lty = 1, col = "black")
```



**Remark 2:**

Vand02 recommend a cut-off value of 0.15 for the pairwise variation. Below this bound the inclusion of an additional housekeeping gene is not required.

### 3 Normalization by geometric averaging

To normalize your data by geometric averaging of multiple reference/housekeeping genes you can proceed as follows

```
> data(SLqPCRdata)
> SLqPCRdata
```

|    | Gene1 | Gene2 | HK1  | HK2  |
|----|-------|-------|------|------|
| A1 | 26.6  | 25.6  | 12.8 | 18.5 |
| A2 | 26.9  | 25.8  | 13.2 | 19.2 |
| A3 | 27.4  | 26.1  | 13.1 | 19.2 |

```

A4  27.7  26.6  13.4  19.5
B1  26.7  25.8  12.9  18.8
B2  24.4  21.5  13.1  18.7
B3  26.5  24.6  12.9  18.7
B4  25.6  23.5  13.8  19.4
C1  28.8  26.6  13.1  19.1
C2  24.4  19.2  13.2  18.5
C3  28.3  25.1  12.9  18.6
C4  25.3  20.6  13.3  19.1
D1  29.3  26.5  12.9  19.0
D2  24.7  18.8  12.7  18.4
D3  27.3  21.1  13.0  18.6
D4  27.3  21.3  13.1  18.4

```

```
> (relData <- apply(SLqPCRdata, 2, relQuantPCR, E = 2))
```

|    | Gene1      | Gene2       | HK1       | HK2       |
|----|------------|-------------|-----------|-----------|
| A1 | 0.21763764 | 0.008974206 | 0.9330330 | 0.9330330 |
| A2 | 0.17677670 | 0.007812500 | 0.7071068 | 0.5743492 |
| A3 | 0.12500000 | 0.006345722 | 0.7578583 | 0.5743492 |
| A4 | 0.10153155 | 0.004487103 | 0.6155722 | 0.4665165 |
| B1 | 0.20306310 | 0.007812500 | 0.8705506 | 0.7578583 |
| B2 | 1.00000000 | 0.153893052 | 0.7578583 | 0.8122524 |
| B3 | 0.23325825 | 0.017948412 | 0.8705506 | 0.8122524 |
| B4 | 0.43527528 | 0.038473263 | 0.4665165 | 0.5000000 |
| C1 | 0.04736614 | 0.004487103 | 0.7578583 | 0.6155722 |
| C2 | 1.00000000 | 0.757858283 | 0.7071068 | 0.9330330 |
| C3 | 0.06698584 | 0.012691444 | 0.8705506 | 0.8705506 |
| C4 | 0.53588673 | 0.287174589 | 0.6597540 | 0.6155722 |
| D1 | 0.03349292 | 0.004809158 | 0.8705506 | 0.6597540 |
| D2 | 0.81225240 | 1.000000000 | 1.0000000 | 1.0000000 |
| D3 | 0.13397168 | 0.203063099 | 0.8122524 | 0.8705506 |
| D4 | 0.13397168 | 0.176776695 | 0.7578583 | 1.0000000 |

```
> geneStabM(relData[, c(3, 4)])
```

|  | HK1       | HK2       |
|--|-----------|-----------|
|  | 0.2574717 | 0.2574717 |

```
> (exprData <- normPCR(SLqPCRdata, c(3, 4)))
```

|    | Gene1    | Gene2    |
|----|----------|----------|
| A1 | 1.728585 | 1.663601 |



A2 1.689720 1.620623  
 A3 1.727684 1.645714  
 A4 1.713602 1.645553  
 B1 1.714500 1.656708  
 B2 1.558954 1.373669  
 B3 1.706201 1.583870  
 B4 1.564586 1.436241  
 C1 1.820707 1.681626  
 C2 1.561410 1.228651  
 C3 1.826986 1.620401  
 C4 1.587369 1.292483  
 D1 1.871526 1.692677  
 D2 1.615795 1.229836  
 D3 1.755636 1.356920  
 D4 1.758402 1.371940

## References

- [1] Jo Vandesompele, Katleen De Preter, Filip Pattyn, Bruce Poppe, Nadine Van Roy, Anne De Paepe and Frank Speleman (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* 2002, 3(7):research0034.1-0034.11 <http://genomebiology.com/2002/3/7/research/0034/> 1