

**GEAR-Symposium, 27.-31.8.2001, Cairo, Egypt**

**“Umweltverschmutzung in Ägypten:  
Folgen für Mensch, Tier und Pflanze”**

## **Studies on the bacterial etiology of pneumonia in camel in northern Jordan**

**Dr. Saeb N. El-Sukhon  
E-Mail : [sukhon@just.edu.jo](mailto:sukhon@just.edu.jo)  
Tel. 00962 2 7248248**

**Department of Basic Veterinary Medical Sciences,  
Faculty of Veterinary Medicine  
Jordan University of Science & Technology  
P.O. Box 3030  
Irbid, Jordan**

### **ABSTRACT**

Pneumonic lungs (71) of 42 camel were examined for their bacterial etiology. The dominant isolated bacteria were identified as *Klebsiella pneumoniae*, *Pasteurella multocida*, *Pasteurella haemolytica*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*. They were determined in 17 (23.9%), 23 (32.4%), 13(18.3%), 9(12.6%) and 4 (5.6%) isolates, respectively. Each of the *Escherichia coli* and the yeast *Candida albicans* were identified twice. The resistance toward the penicillin, amoxicillin, ampicillin, kanamycin, gentamicin, ciprofloxacin, amikacin, enrofloxacin, tetracycline, erythromycin, streptomycin, co-trimoxazol, nitrofurantoin was determined. Fourteen (61%) and 6 (46%) isolates of the *Pasteurella multocida* and the *Pasteurella haemolytica*, respectively, were resistant to the cidal effect of the camel, calf and sheep sera where their count ( $\log_{10}$ ) increased and ranged between 0.2 – 3.6 logs after 1-3 hours interaction time.